



DEEP INELASTIC SCATTERING AND THE EMC EFFECT

by

Gerald Vincent Dunne , B.Sc.(Ma),Hons.

Thesis presented for the degree of Master of Science
in the Department of Physics at The University of Adelaide.

February 1986

AWARDED 28th MAY 1986

Statement:

This thesis contains no material which has been accepted for the award of any other degree or diploma in any University and that, to the best of my knowledge and belief, the thesis contains no material previously published or written by another person, except where due reference is made in the text of the thesis.

I consent to the thesis being made available for photocopying and loan if applicable, provided due acknowledgement is given.

Gerald V. Dunne

Contents

	page
<u>Abstract</u>	(i)
<u>Introduction</u>	(ii)
<u>1. Introduction to Deep Inelastic Scattering</u>	
1.1 Kinematics	1
1.2 Structure Functions	1
1.3 The Deep Inelastic Region	5
<u>2. The Quark-Parton Model</u>	
2.1 Introduction	7
2.2 Infinite momentum frame and Bjorken scaling	7
2.3 Free Quark Model	12
2.4 Sum Rules in the Quark Parton Model	15
2.5 Beyond the Quark-Parton Model	17
<u>3. Operator Product Expansions</u>	
3.1 Introduction	20
3.2 Short distance OPE	21
3.3 Light-cone OPE	25
3.4 Application of the light-cone OPE to DIS	28
<u>4. The Renormalization Group and DIS</u>	
4.1 Anomalous dimensions	35
4.2 Asymptotic freedom	37
4.3 Renormalization Group applied to DIS moments	39
4.4 Sum Rules in QCD	48

5. The EMC Effect

5.1	Introduction	54
5.2	Target dependence in QCD	57
5.3	Change in confinement scale	58
5.4	Space-time picture of DIS; Correlation functions	64
5.5	An Inconsistency	68

6. The Off-Shell Model

6.1	Introduction	70
6.2	Fermi averaging	71
6.3	The off-shell model	75
6.4	Results	78
6.5	The Deuteron	90

Conclusion	95
------------	----

Acknowledgements	96
------------------	----

Appendix A: nucleon structure function parametrization	97
--	----

Appendix B: program for calcium	100
---------------------------------	-----

References	103
------------	-----

Abstract:

This thesis contains a review of theoretical and experimental aspects of deep inelastic scattering. The main predictions of quantum chromodynamics are presented following the introduction of the quark-parton model, operator product expansions and the renormalization group. The recently discovered EMC effect suggests the need for a significant change in the usual QCD treatment of deep inelastic scattering from nuclear targets. The currently popular model involving changes in the quark confinement scale from target to target is discussed, and an inconsistency with the space-time picture of deep inelastic scattering is pointed out. A new explanation, the off-shell model, is proposed and the results for various nuclei are compared with experimental data.

Introduction:

Deep inelastic scattering provides a unique opportunity in the study of the strong interactions because it is both experimentally and theoretically accessible. The agreement between experiment and the predictions of perturbative QCD is quite remarkable. This thesis provides an extensive review of the theoretical predictions of QCD concerning the deep inelastic region. A significant problem with the usual treatment of deep inelastic scattering from nuclear targets has come to light recently. This thesis proposes a new explanation for this problem, the so-called 'EMC effect'.

Chapter 1 contains a review of the basic kinematics of deep inelastic scattering, defining the important kinematic variables, and the 'structure functions'. The second chapter discusses the quark-parton model which was the first serious theoretical treatment of deep inelastic scattering. The main prediction of the quark-parton model is 'Bjorken scaling', a phenomenon observed (approximately) in experiments. The operator product expansion, one of the major mathematical tools in the QCD discussion of deep inelastic scattering, is introduced in chapter 3. Then, in Chapter 4, the field theory quantum chromodynamics is used in conjunction with the operator product expansion and the renormalization group to predict the behaviour of the structure functions as functions of the kinematic variables. Chapter 5 contains a discussion of the recently discovered (1983) EMC effect. The essence of the EMC discovery is that deep inelastic scattering of leptons from nucleons has a significant dependence on the target nucleus. An inconsistency in the currently most popular 'explanation' of the EMC effect is pointed out in this chapter. In the

(iii)

final chapter, a new parameter-free model is proposed which consistently incorporates target dependence into the QCD theory of deep inelastic scattering. This involves accounting for the 'off-shell-ness' of the nucleons in a given nucleus due to nuclear binding effects. This necessitates a modification of the usual Fermi averaging procedures and also introduces a ' Q^2 - rescaling' effect. The new method is applied to many different target nuclei, and good agreement with experimental data is obtained.